

Heritability, correlation and selection response estimates of some traits in fish populations

Gerald W. Friars^{1,2} and Peter J. Smith³

^{1.} 397 Prince of Wales St., St. Andrews, N. B., Canada E5B 1R1

^{2.} Chief Scientist of the Atlantic Salmon Federation's Salmon Genetics Research Program between 1985 and 1995

^{3.} NIWA, Private Bag 14 901, Wellington, New Zealand

Table of Contents

	Page No.
Foreword.....	7
1. Introduction.....	7
2. History of the heritability concept	7
3. Within and between population variation in traits of fish.....	8
4. Estimation of heritability	9
5. Heritability estimates in fish	11
5.1 Comparisons of heritability estimates for traits related to growth and size, age at maturation, and survival	15
5.2 Growth-related traits	15
5.3 Maturation and other age-related traits	16
5.4 Survival-related traits.....	16
6. Related individuals and inbreeding.....	18
7. Multiple Traits	22
7.1 Selection responses	24
7.2 Genetic correlations in fish	25
8. Genotype x environment interactions	25
9. Linking quantitative and qualitative genetics	29
10. Recent references	29
11. Summary.....	30
Acknowledgements.....	31
References.....	31

List of Tables

	Page No.
Table 1 Tests of differences in estimates of heritability (both narrow (N) and broad (B) sense) of growth and size (G) from Figure 1, age at maturation (A) from Figure 2, and survival and survival related traits (S) from Figure 3.....	15
Table 2 Ranks of families at 7°C and 15°C for growth traits in rainbow trout <i>Salmo gairdneri</i>	26
Table 3 Demonstration of log transformation dissipation of an interaction	30
Table 4 Heritability estimates of certain traits in fish in a sample of papers Published in 2009.....	30

List of Figures

	Page No.
Figure 1	12
Figure 2	13
Figure 3	14
Figure 4	19
Figure 5	20
Figure 6	21
Figure 7	23

List of Figures (cont'd.)

	Page No.
Figure 8 Specific growth rates of strains of three trout species and one hybrid reared at two temperatures (with upper 95% confidence intervals). Brook trout <i>Salvelinus fontinalis</i> , lake trout <i>S. namaycush</i> , rainbow trout <i>Salmo gairdneri</i> and the splake hybrid <i>S. namaycush</i> x <i>S. fontinalis</i> . Source Sadler <i>et al.</i> 1986	27
Figure 9 Family rank differences between photoperiod treatments (16 h and natural) for November gains in length and weight, ranked on the basis of increasing initial family means of Atlantic salmon. Source: Hank <i>et al.</i> 1989.....	28

List of Boxes

	Page No.
Box 1 Methods for estimating heritability.....	9
Box 2 Heritability estimates in sea ranching.....	17

Foreword

The material in this dissertation was developed after a presentation at a Fisheries-induced Adaptive Change Workshop coordinated by The International Institute for Applied Systems Analysis, Luxenburg, Austria on March 14-17, 2001, with an intent to publish this dissertation and companion manuscripts in a volume pertaining to that workshop. Although peer review acceptance of this work has been completed, no publication has been forthcoming as of March 2010. Additional commentary on 2009 papers related to this work is included. The paper is now being posted on the Atlantic Salmon Federation website in order to make the information available, and to serve as a benchmark for the rate of advancement of the field.

1. Introduction

Fishing is a major contributor to adult mortality in commercially important fish stocks. Size at harvest and age at maturation, conventionally considered components of life history traits, a term not conventional in animal and plant breeding but used in fisheries, have shown long-term declines in heavily exploited fish stocks (Ricker 1981, Allendorf *et al.* 1987, Law and Grey 1989, Smith 1994). Changes in life history traits result from genetic changes due to selection, both natural and fishery induced and from non-genetic changes due to shifts in the biotic and physical environment. The genetic change or selection response is a product of the selection differential and heritability. Selection intensity can be estimated in wild populations, but there are few estimates of heritability for traits in such populations. Estimates of heritability for key life-history traits are essential for untangling the genetic and environmental components of observed life-history changes in fish stocks.

Heritability estimates have been used extensively in animal breeding programs (Hofer 1998). The high reproductive potential in many fish species can increase selection differentials because few individuals are needed to replace populations. The increasing number of heritability estimates for fish species under aquaculture provides a background for considering fisheries-induced adaptive changes in natural populations. In this chapter the history of the heritability concept is reviewed, before considering heritability estimates in fishes, and the key phenomena of inbreeding, multiple traits, and genotype x environment interactions.

2. History of the heritability concept

An appreciation of the concept of heritability can be gained from its historical development. Concepts of quantitative variation formed the underlying principles of heritability. Knowledge derived from empirical observations on the existence of variability and its random origin was paramount in the emergence of quantitative genetics (Wearden 1979).

Mendelian principles rediscovered in the late nineteenth century, about thirty years after Mendel's publications on peas, introduced complexities in the understanding of quantitative genetics, where discrete classifications of characteristics were not apparent. The merger of the concept of inheritance with that of continuous variation, expressed by Galton (1897) as the regression of offspring on parents, was not without controversy in the scientific community. The hypothesis that several genes contributed to quantitative traits (Yule 1907) was supported by

observations on colour genes in wheat and oats, recorded by Nilsson-Ehle early in the twentieth century (see review by Akerberg 1986). The development of a statistical theory involving correlation between relatives by Fisher (1918) represented a breakthrough in the analyses of quantitative traits, and was enhanced by the publication on systems of mating by Wright (1921). Many of the applications of statistics and matrix algebra in animal breeding stem from the pioneering works of Fisher and Wright (see review by Hofer 1998).

The exact origin of the term “heritability” is obscure, but Wright (1921) used the symbol h^2 to define the proportion of the variance that can be ascribed to heredity in the piebald colour pattern in guinea pigs. The term heritability appears frequently in the animal breeding literature (e.g. Lerner, 1950) where it is defined in a broad sense as “the proportion of the total variance that is directly due to genetic differences”. $h^2 = \sigma^2_G / \sigma^2_P$, where σ^2_G = the genetic variance, and σ^2_P = the phenotypic variance of the stock. Heritability is defined in a narrow sense as the fraction of the genetic variation that is due to genes with a simple additive effect, $h^2 = \sigma^2_A / \sigma^2_P$ (σ^2_A = the additive genetic variance). The realized heritability is defined as R/S , where R (the response to selection) is the difference in average phenotype between the unselected stock and the progeny of the selected parents reared under the same conditions, and S (the selection differential), is the difference in average phenotype between parents and the stock from which they were selected (Ricker 1981). However, environmental conditions are not necessarily the same between generations and estimates of R/S often involve control populations (eg Friars *et al.* 1997).

A literature search for the words “heritability” and “fish” found 148 references in Aquatic Sciences and Fisheries Abstracts (Cambridge Scientific Abstracts in 2000). Growth rate and size, age at maturation, and survival or survival indicator traits were included in the search. Most heritability studies on farmed fish were carried out in the twenty-five years prior to 2001 and in the subsequent nine years through 2009, but the results provide a background for understanding changes in comparable traits in wild fisheries. Developments in molecular biology provide the opportunity for new applications, including heritability estimates and the prediction of genetic gains. The establishment of genetic markers may be feasible in populations mixed during rearing (Ritland 1996).

3. Within and between population variation in traits of fish

The practical estimation of heritability is usually confined to a single population under specific conditions. However more broad heritable variation between species, local populations, stocks and strains also needs to be recognized (see Roff 1992 for an overview).

Variances among river stocks of Atlantic salmon (*Salmo salar*) have been reported for size at return, percent return, day of return, and percent of grilse (Bailey and Saunders 1984). Diallel crosses of four river stocks of salmon revealed larger effects due to the stock of the dams than that of the sires (Friars, Bailey and Saunders 1979). Differences in resistance to the monogenean *Gyrodactylus salaris* were found between populations of Atlantic salmon (Bakke *et al.* 1990) and between populations of charr (Bakke *et al.* 1996). However, strain effects accounted for only 1.8 to 7.4 percent of total variance in size, development, and survival traits in several river stocks of Atlantic salmon (Gjedrem *et al.* 1991a), indicating that heritable variation within stocks is of more consequence.

Mixing stocks from different geographical regions may result in shifting gene frequencies. When a large number of Atlantic salmon escape from sea-cages, enter streams and interbreed with wild stocks, they may disturb adapted gene pools (Crozier 1993). An update on the interaction between aquaculture and wild stocks can be found at ICES (2009).

Molecular techniques have been used to measure genetic variation within and between many fish stocks. However the measurement of simple Mendelian inherited characters, detected with allozymes and molecular DNA variation, is not directly amenable to heritability analyses of quantitative variation, although such techniques are being developed (Ritland 1996).

4. Estimation of heritability

Genetic gain or selection response R is predicted by the product of heritability (h^2) and the selection differential S. Various methods of estimating heritability are outlined in Box 1. Estimates of heritability are generally dependent on the assumptions of large random mating populations in the absence of migration, mutation, and selection, and most methods have been developed for diploids with normal meiosis. Sex-specific differences in recombination may also need to be considered (Sakamoto et al 2000).

BOX 1 Methods for estimating heritability

A. Regression between parents and offspring

Regression methods estimate the extent of the transmission of a trait from parent to offspring: $y = a + bx$ where x and y are the mean values of the trait for the parents and offspring respectively, a is the intercept, and b the regression coefficient. When the mean values of a trait, derived from both parents, is used the heritability is equivalent to the regression coefficient, $h^2 = b$, but if the offspring are compared with just one parent the heritability, $h^2 = 2b$.

B. Hierarchical matings

The following model describes the sources of variation in the population:

$Y_{ijk} = \mu + S_i + D_{ij} + e_{ijk}$ where Y_{ijk} is the measurement of a trait on the k^{th} progeny of the j^{th} dam mated to the i^{th} sire, μ is the overall mean of the population, S_i is the effect of the i^{th} sire, D_{ij} is the effect of the j^{th} dam mated to the i^{th} sire, and e_{ijk} is the environmental plus genetic segregation effect on the k^{th} progeny of the j^{th} dam mated to the i^{th} sire.

Heritability is estimated from the sire component (h^2_s), the dam component (h^2_D), and the sire and dam component (h^2_{S+D}):

$$h^2_s = 4 \sigma^2_s / (\sigma^2_s + \sigma^2_D + \sigma^2_e);$$

$$h^2_D = 4 \sigma^2_D / (\sigma^2_s + \sigma^2_D + \sigma^2_e); \text{ and}$$

$$h^2_{S+D} = 2(\sigma^2_s + \sigma^2_D) / (\sigma^2_s + \sigma^2_D + \sigma^2_e).$$

BOX 1 cont'd.

where σ^2_S is the sire component of variance, σ^2_D the dam component of variance, and σ^2_e the environmental plus genetic segregation component of variance. Dam components are inflated by maternal effects, dominance and epistatic variance to a greater extent than sire components where dams are nested within sires. Also where fertilization is in vitro, as in fish, sires can be nested within dams.

C. Factorial matings

Sets of matings where sires from each strain are mated with dams of each strain allow for partitions of variance components where the theoretical expectations of additive and non-additive effects are as below (McKay *et al.* 1986):

- σ^2_A = variance due to additive genetic effects;
- σ^2_D = variance due to dominance;
- $\sigma^2_{AA}, \sigma^2_{AD}, \sigma^2_{ADD}$ = variance due to epistatic interactions of additive-additive, additive-dominance, and dominance-dominance types respectively;
- σ^2_{Ec} = variance due to environmental effects between tanks;
- σ^2_{EG} = other environmental effects not due to tank effects;
- σ^2_M = variance due to maternal effects.

Variance Component	σ^2_A	σ^2_A	σ^2_{AA}	σ^2_{AD}	σ^2_{DD}	σ^2_{Ec}	σ^2_{EG}	σ^2_M
Between Families	1/2	1/4	1/4	1/8	1/16	1	0	1
Within Families	1/2	3/4	3/4	7/8	15/16	0	1	0

D. Breeding values

The prediction of breeding values in large data sets has been used extensively in animal breeding. The selection criterion has been based on best linear unbiased prediction (BLUP) of additive genetic effects. BLUP requires knowledge about variance components that have been estimated in practice. Restricted maximum likelihood (REML) has become the method of choice for the estimation of variance components of mixed linear models in animal breeding (Hofer 1998).

E. Inbred lines

The use of variance between and within inbred lines can be used to estimate genetic variance and heritability $h^2_t = h^2_0(1-F_t)/(1-h^2_0 F_t)$

where h^2_0 is the heritability in the base population, h^2_t and F_t are the heritability and inbreeding coefficients at time represented by generations t ,

F. Linear heritability estimates

Linear estimates are based on selecting prospective parents for which estimates of average genotypic and phenotypic values are obtained from a pedigreed population (Abplanalp 1961). A regression of genotypic on phenotypic values is determined from a straight line fitted through points representing the population mean genotype and phenotype on one hand and the mean genotype and phenotype of selected parents on the other.

G. Clonal lines

Taniguchi *et al.* (1996) produced mitotic gynogenetic diploids, reared them in a common environment, identified lines using DNA fingerprinting and employed a human twin model to separate genetic and environmental variances.

H. Discrete traits

Traits such as mature versus non-mature fall into discrete binomial distributions. Heritability can be estimated from an underlying continuous scale using techniques, which provide a quantification of multiple gene effects (Robertson and Lerner 1949). Also, family means can be considered for such traits, the heritability of which is outlined later in this chapter.

I. Animal model

Personal communication with L.R. Schaeffer resulted in a quotation as follows:

“An animal model incorporates the animal’s records, its parents, and its progeny into one overall estimate of the breeding value of each animal. Pedigree information has to be accurate and complete (personal communication with L.R. Schaeffer). Details can be found at:

www.aps.uoguelph.ca/~Irs/ABModels/ or www.aps.uoguelph.ca/~Irs/ABMethods/”

5. Heritability estimates in fish

A large number of heritability estimates has been made for fish and shellfish - 19 traits in 26 species. Most heritability estimates are for carp and salmonids, which is not surprising given the long history of carp in culture and the more than 100 years of experience with salmonids in enhancement programs.

Three life-history traits in fisheries, growth rate, size at maturation, and survival, were selected from the 19 traits noted in the literature. These traits are relevant to the changes observed in exploited populations of decreases in size and age at maturation, and survival (which bears on the former two traits with respect to the shifts in genetic make-up that relate to the fitness of the stocks). Heritability estimates for the three traits are summarized in Figures 1, 2 and 3; the numbers in the columns relate to the reference list at the end of this chapter. The histograms show wide distributions for each class of traits.




Figure 1. Estimates of heritability of growth, size and related traits in several species of fish. Narrow sense heritabilities (NSH) contain principally additive genetic effects, including realized heritabilities. Broad sense heritabilities (BSH) contain both additive and non-additive genetic effects. Numbers in the histograms represent points that can be traced to references where the respective papers are denoted in bold superscript in front of the authors. The information predates 2002.




Figure 2. Estimates of heritability of age at maturation, age at smolting and traits related to reproduction in several species of fish. Narrow sense heritabilities (NSH) contain principally additive genetic effects, including realized heritabilities. Broad sense heritabilities (BSH) contain both additive and non-additive genetic effects. Numbers in the histograms represent points that can be traced to references where the respective papers are denoted in bold superscript in front of the authors. The information predates 2002.




Figure 3. Estimates of heritability of survival, disease resistance and related traits in several species of fish. Narrow sense heritabilities (NSH) contain principally additive genetic effects, including realized heritabilities. Broad sense heritabilities (BSH) contain both additive and non-additive genetic effects. Numbers in the histograms represent points that can be traced to references where the respective papers are denoted in bold superscript in front of the authors. The information predates 2002.

5.1 Comparisons of heritability estimates for traits related to growth and size, age at maturation, and survival

There are more references for growth and size than for age at maturation or survival- related traits (Figures 1, 2, and 3). The wide range in heritability estimates for these traits emphasizes the fact that inferences are to populations with a specific genetic make-up in a specific environment. Small sample sizes and biases in estimation procedures also contribute to the wide distributions of estimates. Tests of differences between medians (Table 1) were performed using procedures outlined by Siegel (1956). The probabilities of type 1 errors (rejecting the null hypothesis when true) are generally high, with the exception of the test for narrow- versus broad-sense heritability estimates for growth related traits, because of the large number of estimates. Consequently, statements about the relative levels of heritability of traits must be interpreted with caution. Furthermore, expression of quantitative traits can be highly dependent upon the environment and is likely to be more complex under natural than experimental conditions (Ritland 1996, Merilä and Sheldon 2000, Hermida *et al.* 2002). Thus heritabilities of traits in wild fish populations are needed for estimating the selective effects of fisheries. While few data exist for wild populations, with the exception of returning salmonids (Jonasson, 1997), the use of molecular markers to infer relatedness may provide a tool for studying quantitative variation in natural populations (Ritland 1996).

Table 1. Tests of differences in estimates of heritability (both narrow (N) and broad (B) sense) of growth and size (G) from Figure 1, age at maturation (A) from Figure 2, and survival and survival related traits (S) from Figure 3.

	Trait					
	G	G	A	A	S	S
	N	B	N	B	N	B
Number of estimates	120	75	20	8	49	48
Median	0.28	0.43	0.51	0.21	0.16	0.28
Contrast*	GN-GB	AN-AB	SA-SB	GN-AN	GN-SN	AN-SN
χ^2	8.33	0.04	0.51	1.00	2.10	1.20
P type 1 error	<0.01	<0.95	<0.50	<0.28	<0.20	<0.20

*The contrasts are between narrow and broad sense heritability estimates and between traits.

5.2 Growth-related traits

Most estimates for growth related traits fell within the range 0.10 to 0.60 for narrow sense heritabilities (Fig 1); the spikes in frequencies below 0.175 are mostly for salmonids in sea cages, and may indicate lower heritabilities for marine than for land-based aquaculture populations. The heritabilities in the upper Figure 1, include maternal effects and non-additive genetic effects, namely dominance and epistasis, and generally result in a slightly higher mean than that for only additive effects. The lower estimates contain largely additive effects together with minor amounts of epistasis. The significant difference ($P < 0.01$; Table 1) reflects this trend where the heritability estimates (Figure 1) have a higher median value in the broad sense than

those in the narrow sense. Nevertheless, these contributions pertain to population phenomena that may have biological significance.

5.3 Maturation and other age-related traits

There is a wide distribution of heritability estimates of age at maturation related traits (Figure 2). Some of the estimates are for traits such as the proportion of smolts at certain ages, and others for proportions exhibiting sexual maturity at specified ages. In general, heritabilities for age at maturation traits (Figure 2) do not appear to be different to values for growth-related traits (Table 1). The limited number of estimates available from the literature indicates that considerable responses could be realized through selection for age at maturation. If fishery- selection acts on age-related traits, it is reasonable to expect that changes in age at maturation will occur.

Age at maturation may be related to fitness, based on the relative numbers of offspring surviving from fish maturing at different ages. For instance, Atlantic salmon grilse reproduce one year before two-sea-winter contemporaries, but produce fewer eggs – influencing the number of offspring per year. Consequently the lower heritability of reproductive traits compared with growth traits, due to higher components of environmental variance, may be limited to specific situations (Merilä and Sheldon 2000).

A review of the early literature on sea-age and maturation in Atlantic salmon concluded that heritable factors were involved in maturation after one-sea-winter (grilse) versus two-sea-winters (Gardner 1976). That view is justified where heritability estimates of non-grilse salmon were of 0.19, 0.21 and 0.22 in three populations, based on fullsibs (O’Flynn *et al.* 1992). Also, a heritability estimate of 0.48 was found for a similar trait in rainbow trout by Gjerde (1988), using crosses of parents from different ages at maturation. The findings of Ritter *et al.* (1986), working with wild stock, agree with these results where the phenomenon of 1-sea-winter versus older salmon representing age at maturation is heritable. The wide range of estimates (Figure 2) substantiates the heritable nature of age at maturation across a number of species.

5.4 Survival-related traits

Traits relating to survival generally have lower heritabilities than growth rate and age at maturation (Figure 3 and Table 1), possibly due to the complexity of survival traits that include disease resistance and physiological phenomena. However, selection for disease resistance has been successful in developing resistance to furunculosis in Atlantic salmon (Bailey *et al.* 1993). A heritability of 0.24 for swimming stamina, a trait possibly related to survival, was reported in Atlantic salmon (Hurley and Schom 1984), and similar results have been found for the three-spine stickleback (*Gasterosteus aculeatus*), (Garenc *et al.* 1998).

Behavioral traits, such as schooling and migratory tendencies (Rose 1993), could affect survival in a fishery. In Atlantic cod, there are differences in migratory behaviours between coastal and north-east Arctic cod (Godo 1995) and differences in activity patterns between cod stocks in the Irish and North Seas (Righton *et al.* 2001). If fisheries impose selective forces on such traits, shifts in characteristics such as growth and/or age at maturation will be realized due to genetic dependencies between traits. These changes will depend not only on

direct selection on the traits, but will be influenced by genetic correlations among behavioral and other traits.

Return rates of Atlantic salmon, released to sea, represent an aspect of survival. Bailey and Saunders (1984) reported differences in the average date of returns between strains of Atlantic salmon. A summary of some of the heritability estimates of return rates of Atlantic salmon from the findings of Jonasson and colleagues is depicted in Box 2. A genetic equilibrium aspect of return rates was reported in Atlantic salmon where such rates for two synthetic gene pools, produced from crosses of river stocks, revealed percentages of 0.60, 0.18 and 0.43 for the first pool, and 0.68, 0.06 and 1.00 for the second pool, over three successive generations (Friars *et al.* 1994), emphasizing the problem when new genes are introduced into populations. In addition, the return rates of the two synthetic gene pools increased after random mating, at a time when many neighbouring Bay of Fundy rivers were experiencing decreased return rates (Friars *et al.* 1994).

BOX 2 Heritability estimates in sea ranching

Sea ranching is an aquaculture operation that uses the homing habits of anadromous fish, in particular, salmon. The juvenile freshwater stages are raised under controlled hatchery conditions, and at smoltification (the physiological changes prior to entering sea water) the salmon are released to sea where they feed as wild fish. After 1-3 years at sea, the sexually mature fish return to the release site and are caught before they can spawn. Fish returning after one year are known as grilse and after two years as two-sea-winter salmon.

Higher estimates of heritability (Box 1) and of weight and survival using dam (female) as opposed to sire (male) components of variance, have been reported in sea-ranched Atlantic salmon. These are consistent with the estimates in Figure 1 for weight and length with respect to narrow-sense versus broad-sense heritability estimates but not for age at maturation (Figure 2) where the contrast (i.e., the statistical test of differences) is nonsignificant (Table 1). The higher estimates of the heritability of weight in farmed as opposed to sea-ranched fish are not significantly difference in light of the means and standard errors of sea-ranched and farmed grilse (Table in this box; Jonasson and Gjedrem 1997). It has been assumed that environmental variances are higher in wild stocks than in farmed stocks, partly because environmental effects inflate the denominators of heritability estimates and partly because of the higher standard deviations and of factors other than additive genetic effects (Jonasson and Gjedrem 1997).

Heritability estimates in Atlantic salmon				
<u>Traits</u>	<u>Sire Component Of variance^a</u>	<u>Dam component of variance^a</u>	<u>Animal model^b</u>	<u>Reference</u>
Freshwater (12 weeks after first feeding)				
Survival	0.04±0.04	0.34±0.04		Jonasson 1993
Weight	0.16±0.02	0.36±0.03		Jonasson 1993
Length	0.10±0.02	0.39±0.03		Jonasson 1993
Sea-ranched grilse				
Weight (1989)			0.20±0.10	Jonasson and Gjedrem 1997 ^a

Weight (1991)			0.23±0.12	Jonasson and Gjedrem 1997 ^a
Weight (land-based)			0.31±0.11	Jonasson and Gjedrem
Farmed grilse				
Weight (1989)			0.31±0.17	Jonasson 1997
Weight (1991)			0.27±0.10	Jonasson 1997
Return rate				
Grilse	0.12	0.12		Jonasson <i>et al.</i> 1997 ^a
2-sea-winter	0.02	0.11		Jonasson <i>et al.</i> 1997 ^a

Correlation estimates in Atlantic salmon

Traits	Phenotypic	Genetic	Reference
Eggsize			
Survival	0.23**		Jonasson 1993 ^a
Weight	0.233		Jonasson 1993 ^a
Length	0.22**		Jonasson 1993 ^a
Survival			
Weight		0.31±0.26	Jonasson 1993
Length		0.39±0.26	Jonasson 1993
Weight-length		0.98±0.01	Jonasson 1993
Weight (relatives in sea-ranched And farmed environments)			
1989	0.14	0.42±0.28	Jonasson 1997
1991	0.09	0.27±0.34	Jonasson 1997

^aWhere dams are nested within sires in the mating system, the dam components of variance contain maternal effects and more nonadditive genetic effects than the sire components of variance.

^bAn animal model optimally combines information on the individual, its parents, sibs, and progeny to estimate the additive genetic merit of each individual. This information allows the estimation of narrow-sense heritability.

**Significance (P) < 0.01.

Other references on Jonasson's work in this area are as follows: Jonasson, J. 1996; Jonasson, J. and T. Gjedrem 1997.

6. Related individuals and inbreeding

Related individuals have genes that are common by descent (Figure 4). This shared inheritance creates correlations between relatives as shown by Fisher (1918). Heritabilities of groups of relatives, such as full or half-sibs, can be estimated by making use of suitable mathematical relationships, and applied in the development of selection and mating programs. Information on relatives has been used extensively in animal breeding:

1. To increase the accuracy of prediction of the breeding value of individuals. An example of this is shown in terms of information from sibs (Figure 5), adapted from Lerner (1958) using an application of path coefficients by Wright, (1921).
2. To establish the breeding value of groups of relatives. One example of the heritability of a group of relatives is depicted in Figure 5. The increased heritability of family means over individual observations is depicted in Figure 6. Caution is needed where the use of information from relatives tends to increase inbreeding during selection.

The use of information on relatives may develop in fisheries as genetic tagging technology evolves (e.g. Herbinger *et al.* 1995; Hansen *et al.* 2001).




Figure 4. The biometric basis of family selection, adapted from Lerner (1958) using an application of path coefficients by Wright (1921).

A represents the genetic (r^G) and phenotypic (r^P) relationships between n members of a family. The genetic values of each individual are represented by G_1 to G_n ; the family average is represented by \bar{G} . Individual phenotypic values are represented by P_1 to P_n and the family average by \bar{P} .

The path between genotype and phenotype (square root of heritability) is indicated by h .
 I = selection index, p (with subscripts) = path coefficients and r (with subscripts) = correlation coefficients.

The values of the paths g and p are:
 $g = 1/\sqrt{n[1+(n-1)r^G]}$ $p = 1/\sqrt{n[1+(n-1)r^P]}$




Figure 5. Heritability of a family average. The heritability of a family average h_{fa}^2 is analogous to $r_{\bar{A}\bar{P}}^2$. Adapted from Lerner 1958.




Figure 6. Ratio of family to individual heritability. The curves are based on the assumption that the relationship between full sibs (r^G) is 0.5 and that $r^p = r^G h^2$. (h_f^2 = heritability of family averages). The ratio h_f^2/h^2 depends on heritability and on number in a family of full-sibs. Adapted from Lerner 1958.

In aquaculture inbreeding has led to a decrease in survival and growth related traits in farmed rainbow trout (Gjerde *et al.* 1983) and catfish (Smitherman *et al.* 1996) and similarly in mitotic-gynogenetic diploid fish (Tsuji-mura and Taniguchi 1996). The development of inbred lines of fish has potential in aquaculture. Heterosis was realized in the crosses of inbred lines of rainbow trout, but the costs and delays in developing inbred lines did not justify the heterotic gains (Gjerde 1988). Large population sizes (N) and effective population sizes (N_e) indicate that inbreeding is not likely to be a problem in most marine fish populations, or in broad samples from the wild for aquaculture.

Bi-directional selection can reveal different rates of progress. For example inbreeding increased the realized heritability more in the downward than in the upward direction in the flour beetle *Tribolium castaneum* (Jui and Friars 1974; McLeod and Friars 1986). In agriculture, goals are usually directed towards increased yields through selection in the upward direction. Conversely fisheries, that preferentially capture larger fish, are directing downward responses in natural populations.

7. Multiple Traits

Several traits such as growth or size, age at maturation, and survival need to be considered simultaneously in fish populations. Heritability estimates pertain to individual traits, and models of multiple traits require the integration of heritabilities with correlations between traits. The yield from a population is influenced by a number of traits, especially more tangible components such as those depicted in Figures 1, 2, and 3. Therefore, in addition to heritability, the genetic, environmental, and phenotypic correlations have to be considered. Figure 7 shows the interrelationships of these phenomena, as they relate to an aggregate or multiple genotype and ultimately an index or collective phenotype.




Figure 7. The genetic (G), phenotypic, and environmental (E) correlations between traits A and B. Adapted from Lerner 1958.

The path diagram (Figure 7) depicts how multiple traits can be combined to form a hypothetical trait in the form of an index. Three types of effects, genetic, environmental, and economic weights, bear on the aggregate index scale. The phenotypic correlation (r_p) can be defined as the sum of the genetic path ($h_A r_A h_B$) and the environmental path ($e_A r_E e_B$). The relative economic weights assigned to each trait affect the index to which directional selection can be applied, and are used to provide profit criteria for each trait. In the case of fisheries, size probably receives more emphasis than other traits, such as age at maturation and survival.

An index was developed for Atlantic salmon with a view to increasing the proportion of smolts, developing from hatchery-reared eggs, after 18 months in fresh water and the market size after approximately 70 weeks in seawater, while decreasing the proportion of grilse (O'Flynn *et al.* 1992). Such methodology is generally considered to be more efficient than the use of independent culling levels on each trait. Progress from selection using an index that involved market size and non-grilse was realized in Atlantic salmon by Friars *et al.* (1993) where size at market age was increased, while the incidence of grilse was reduced. A reversal of this selection

result appears to take place where large fish are harvested in commercial fisheries. Consequently, earlier sexual maturity and smaller size results (Smith 1994).

7.1 Selection responses

The selection response on a trait is determined by the product of the narrow sense heritability and the selection differential. The prediction of such response depends on the reliability of heritability estimates where genetic parameters are subject to change under selection. For instance, Clayton *et al.* (1957a) found fair agreement between early realized selection gains and gains predicted via an estimate of heritability and selection differentials, whereas predictions were poor in the long-term for bristle number in *Drosophila* (Clayton *et al.* 1957b). However, Enfield (1977) realized 26 genetic standard deviations and 15 phenotypic standard deviations of gain in long-term selection for pupal weight in *Tribolium castaneum*, suggesting that new sources of variation accrue to inhibit the exhaustion of genetic variance. Examples of long-term selection experiments have not been well documented in fish. However, Hershberger *et al.* (1990) found that heritabilities for weight in coho salmon remained high after four generations of selection.

Eight generations of selection in chickens revealed mostly insignificant changes in heritabilities, but changes in genetic correlations were apparent in a study by Friars *et al.* (1962). Larval weight and pupa weight in *Tribolium castaneum*, which would be expected to involve part-whole correlations, showed a positive correlation of 0.55 ± 0.12 and exhibited no detectable change over nine generations of divergent selection (Bell and Burris 1973). Conversely, realized genetic correlations between larva weight and offspring number in *T. castaneum* were positive and decreasing under selection in a wet environment, but in a dry environment, the estimates were near zero and increasing slightly positively under selection (Friars *et al.* 1973). In an optimal environment, such as the wet condition in the *Tribolium* experiments, genetic covariance that positively affects weight and offspring number gets used up under selection. Conversely, depression of the traits in the dry environment probably inhibits the expression of genes that affect both traits positively. Inferences to such a combination of effects could be investigated in a fishery, using data from long-term selection.

Studies by McMillan *et al.* (1995) illustrated that changes in parameters over generations of selection can affect outcomes. Under some simulated conditions, the use of initial genetic parameter values without change could have potentially negative effects on overall genetic gain. Hence, estimation of genetic parameters at varying stages of multiple objective selections is required to attain desired goals. Where selection is being exerted in a fishery, updated estimates of genetic parameters are needed to make decisions related to the multiple traits involved, particularly where evidence in other organisms indicates that changes take place under selection.

A recent high-low controlled selection experiment in Atlantic silversides (*Menidia menidia*) by Conover *et al.* (2009) revealed that “large harvested populations initially produced the highest catch but quickly evolved a lower yield than controls. Small harvested populations did the reverse.” In light of these results, Conover *et al.* recommended that management of fishery stocks need reconsideration.

7.2 Genetic correlations in fish

Genetic correlations result from pleiotropic effects of genes on multiple traits or from chromosomal linkage of genes affecting different traits. Estimates of genetic correlations between traits have been made for several fishes. Eknath *et al.* (1998) found a full-sib estimate of 0.2 for the genetic correlation between survival and market weight of farmed *Tilapia*. Estimates between different measures of growth and size, were found to be universally positive, from swim-up to four years of age in rainbow trout (McKay *et al.* 1986). Similarly, correlations between measures of growth and smoltification, in fresh water, related to measures of growth in seawater in Atlantic salmon were generally positive, but became progressively weaker the longer the fish were in sea water (Friars *et al.* 1990).

Herbinger and Friars (1991) found a strong positive correlation, between condition factor and total lipid content (an indicator of energy reserves) in immature salmonids. Another study on the same stock revealed a positive relationship, between July and November parr length, but a negative association, between July parr length and incidence of presumptive smolts in February (Bailey and Friars 1994).

Relationships between body weight and age at maturation were reported in rainbow trout, with genetic correlations of 0.67 and 0.58 (Crandell and Gall 1993a). Selection for market size, and related traits, after two sea winters in Atlantic salmon, was associated with a lower proportion of fish maturing after one sea winter (Friars *et al.* 1993). Considering the selection for smaller sized fish to reproduce in a natural fishery, these results would suggest a trend for earlier age at maturation in natural populations where larger fish are harvested.

Survival also relates to reproduction rate where fish die before reproducing. For Atlantic salmon, river stocks differed in their proportion of grilse versus multiple-sea-winter salmon, as noted by Bailey and Saunders (1984). If lower survival is experienced by multiple-sea-winter fish, due to longer exposure at sea, an advantage in reproduction countered by fewer eggs per female is experienced by grilse. Hence, counteracting forces can affect the reproductive output due to later age at maturation versus survival to reproduction at a younger age.

Age at maturation has a wide range of heritability estimates (Figure 2) and if correlated with size, as found in Atlantic salmon and rainbow trout, and then fisheries that selectively harvest larger fish will not only reduce size, due to its heritability, but also decrease age at maturation.

8. Genotype x environment interactions

The differential reaction of genotypes, across different environments, is of concern to agriculturists and aquaculturists. Superior performance of one genotype may not be consistent when genotype x environment interactions occur. Selection for genotypes in a hatchery environment may not produce the optimum genotypes for a production environment. Parallel problems could exist in fisheries where shifts in environments, due to natural causes, could cause similar problems in wild fisheries.

In salmonids, strains reacted differently under two temperature treatments (Figure 8). At the chromosomal level, family differences in relative growth of diploid and triploid Atlantic salmon represented interactions with respect to growth responses, which imply that different genes for growth are involved in diploid and triploid backgrounds (Friars *et al.* 2001). Likewise McKay (1982) observed that families changed rank for gain in fork length and instantaneous growth rate for weight (Table 2). Similar family- environment, but not strain-environment effects were observed in Atlantic salmon (Wild *et al.* 1994, Winkelman & Peterson 1994a). The results of Jonassen (1996) with ranched Atlantic salmon were similar for weight, where no significant effect was found for release-site x strain interaction. However, such effects were significant for return rate in two out of three year classes and one out of three for ratio of grilse. Eknath *et al.* (1998), following an extensive study with several strains and farming regimes with Tilapia, also concluded that specialized strains may not be necessary for farming systems in the Philippines.

Table 2. Ranks of families at 7°C and 15°C for growth traits in rainbow trout *Salmo gairdneri*. Source: McKay (1982).

Family ^a	Gain in fork-length				Instantaneous growth rate for weight			
	Period 1 7°C		Period 2 15°C		Period 1 7°C		Period 2 15°C	
1	1	8	4	8	3	13	16	18
2	2	12	17	19	1	5	2	31
3	3	11	3	2	2	8	23	19
4	4	30	15	3	6	28	5	10
5	5	7	1	1	4	18	7	6
(Families 6 to 30 not shown)								
31	31	21	35	35	29	9	32	33
32	32	14	29	22	33	24	24	15
33	33	28	23	15	25	23	25	8
34	34	27	28	12	30	20	15	5
35	35	35	22	31	34	35	31	35

^a Families arbitrarily numbered based on their performance at 7°C in period 1.




Figure 8. Specific growth rates of strains of three trout species and one hybrid reared at two temperatures (with upper 95% confidence intervals). Brook trout *Salvelinus fontinalis*, lake trout *S. namaycush*, rainbow trout *Salmo gairdneri* and the splake hybrid *S. namaycush* x *S. fontinalis*. Source Sadler *et al.* 1986.

Juvenile Atlantic salmon reared under two photoperiods, natural day length and 16 hours, revealed that families reacted differently to the shift in the light regimes (Hanke *et al.* 1989, and Figure 9). These examples are related to productivity in aquaculture but suggest that similar phenomena may be acting in natural populations of fish.




Figure 9. Family rank differences between photoperiod treatments (16 h and natural) for November gains in length and weight, ranked on the basis of increasing initial family means of Atlantic salmon. Source: Hank *et al.* 1989.

The extension of genotype x environment interactions to include several environments has been of concern with terrestrial species. A review of the concepts and biometry of the case of genotype x environment interaction was published by Yamada (1962). A trait, measured in two or more environments, was considered to represent two or more traits and entailed genetic correlations of such traits. That work used random and mixed statistical models. This view of the area relates to the concept of the reaction norm used in ecology and is illustrated in Stearns (1992). Where traits are heritable, as in the case of size at harvest and age at maturation (Figures 1 and 2) the adaptability of gene pools to different environments, including levels of fishing pressure, is relevant to concerns with wild populations, as discussed by Ritter *et al.* (1986).

Interactions are subject to scale. A hypothetical example (Table 3) demonstrates that a log transformation of data can remove an interaction that appears on the untransformed scale. However, the results on a transformed scale often have little practical use. Many of the studies on fish deal with transformed data and do not consider this implication. Observed genotype by environment interactions must be taken into account to be of use in fisheries, where specific gene pools are required for specific environments.

Table 3. Demonstration of log transformation dissipation of an interaction.

Strain	Untransformed weights (gms)		Ln. of weights	
	Environment		Environment	
	E ₁	E ₂	E ₁	E ₂
A	10	20	2.3	3.0
B	15	30	2.7	3.4

9. Linking quantitative and qualitative genetics

Molecular or qualitative markers are generally poor indicators of heritable variation in quantitative traits (Mackay and Latta 2002) and their major application in fisheries has been to define genetic population structure. Application of highly variable microsatellite and minisatellite fingerprints of fish enables stock and pedigreed families to be reared in communal ponds and subsequently traced to individual parents (e.g. Herbinger *et al.* 1995; Vandeputte *et al.* 2001). Such techniques have the potential to be extended to natural populations to infer pedigrees allowing the estimation of genetic parameters (Ritland 1996). The construction of detailed genetic maps with molecular markers allows the identification of chromosomal regions influencing quantitative traits in fish (e.g. Sakamoto *et al.* 1999; Peichei *et al.* 2001). Construction of linkage maps based on a large number of DNA markers is necessary to identify quantitative trait loci (QTL) controlling traits of importance in fisheries. Microsatellite markers, which are widely distributed throughout the genome, are ideal for genome mapping and identification of QTL, and are being applied in salmonids (Sakomoto *et al.* 1999, 2000). The application of this technology to other fishes under aquaculture, such as cod and plaice, will allow the identification of QTL controlling traits important in fisheries, and provide new tools for measuring genetic change in natural populations.

10. Recent references

In 2001, a computer search of the literature revealed 148 papers dealing with heritability in fish. A similar search in 2009, covering the period 2000 – 2009, listed 256 papers. A sample of papers in 2009 indicated that heritability studies have been conducted on numerous species of fish dealing with various traits (Table 4).

Table 4. Heritability estimates of certain traits in fish in a sample of papers published in 2009.

Authors	Species	Traits and heritability estimates
Antonello, J. <i>et al.</i>	Gilthead sea bream (<i>Sparus aurata</i> L.)	Body length post mortem $.38 \pm .07$ pasteurellosis resistance; survival post challenge $.12 \pm .04$ and $.45 \pm .04$ to $.18 \pm .08$ binary dead or alive at specific day.
Bardon, A. <i>et al.</i>	European sea bass (<i>Dicentrarchus labrax</i>)	Spinal deformities – global $.21 \pm .04$; lordosis; $.13 \pm .04$ scoliosis $.13 \pm .04$.
Gheyas, A.A. <i>et al.</i>	Silver carp	Harvest weight $.67$ (.42 – .93 conf. lim.)
Guy, D.R. <i>et al.</i>	Atlantic salmon (<i>Salmo salar</i>)	Infectious pancreatic necrosis percent mortality $.07$ to $.56 \pm <.04$.
Khaw, H.L. <i>et al.</i>	Nile tilapia (<i>Oreochromis niloticus</i>)	Harvest weight $.15 - .41 \pm .04$
Navarro, A. <i>et al.</i>	Gilthead seabream (<i>Sparus auratus</i> L.)	Harvest weight $.34 \pm .06$; fork length $.33 \pm .07$; dressing percentage $.31 \pm .07$; fillet percentage $.12 \pm .04$.
Taylor R.S. <i>et al.</i>	Atlantic salmon (<i>Salmo salar</i> L.)	Amoebic gill disease – gill score observed scale $.23$ to $.48$; survival challenge $.40$ to $.49$; weight $.28$ to $.27$; length $.27$ to $.35$.

Various methods of estimation, including the animal model were employed in these studies (Table 4). Molecular markers were used in some studies to establish pedigrees – a technique that circumvents the confounding of family and environmental effects such as tank effects, as frequently encountered in older studies. In some of these studies, genetic correlations between traits were also estimated.

11. Summary

The concept of heritability, generally defined as the proportion of phenotypic variation due to genetics, has been widely applied in agricultural fields. Extensions of such developments to fish have been studied principally through aquaculture. A survey of heritability estimates on growth, age at maturation, and survival, and inferences to fisheries were considered in this chapter. Summaries of heritabilities indicate that survival is less heritable than growth or age at maturation even though the probabilities are not strong (Table 1). Consideration of correlations

between traits suggest that multiple traits need simultaneous attention in fisheries. The use of molecular techniques may allow estimations of heritabilities and genetic correlations in wild populations in the near future.

Acknowledgements

We are grateful to the European Science Foundation for support; to the University of Guelph, the Department of Fisheries and Oceans (Canada), the Huntsman Marine Science Center, and the Atlantic Salmon Federation for resources; and to Ian McMillan, Laura McKay and Margaret Quinton (University of Guelph), Alex Hanke (Fundy Data Acquisition Ltd., Chamcook, NB) and Bob McDowall (National Institute of Water and Atmospheric Research Ltd = NIWA), and Jarle Mork (Norwegian University of Science and Technology, Trondheim, Norway) for comments on the draft manuscript. Peter Smith acknowledges support from NIWA.

The clerical help of Ellen Merrill of the Atlantic Salmon Federation has been extensive.

References

(Note: the superscripted numbers at the front of some references relate to Figures 1, 2, and 3 for heritability estimates).

Abplanalp, H. 1961. Linear heritability estimates. *Genetical Research* 2: 439.

Åkerberg, E. 1986. A historical survey of breeding research. In: Svalof Editor – G. Olsson 1886-1986, pp 9-34 Publisher –LTs forlag Stockholm.

Allendorf, F.W., Ryman, N., Utter, F. 1987. Genetic and fisheries management: past, present and future. In *Population Genetics and fisheries management* Eds Ryman, N and Utter, F.M. pp 1-20. University of Washington Press, Seattle.

⁰¹Anon. 1996. Resistance to Bacterial Kidney Disease in New Brunswick Farmed Atlantic salmon. A joint project between the Salmon Genetics Research Program, Atlantic Salmon Federation, St.Andrews, NB and the Research and Productivity Council, Fredericton, NB. Canada. 28pp.

Antonello, J., Massault, C., Franch, R., Haley, C., Pellizzari, C. Bovo, G., Patarnello, T., de Koning, D.J., Bargelloni, L.S.O. 2009. Estimates of heritability and genetic correlation for body length and resistance to fish pasteurellosis in the gilthead sea bream (*Sparus aurata* L.). *Aquaculture* 298(1-2): 29-35.

Bailey, J.K. and R.L. Saunders. 1984. Returns of three year-classes of sea-ranched Atlantic Salmon of various river strains and strain crosses. *Aquaculture* 41:259-270

⁰²Bailey, J.K. and Friars, G.W. 1990. Inheritance of age at smolting in hatchery-reared Atlantic salmon (*Salmo salar*). *Genetics in Aquaculture III, Aquaculture*. 85:317 (abstract).

⁰³Bailey, J.K., O'Flynn, F.M. and Friars, G.W. 1991. Abstract. Atlantic Salmon Federation, St. Andrews, NB. Genetic and phenotypic correlations among freshwater growth traits in Atlantic salmon. Aquaculture Association of Canada Meeting June 10, 1991.

Bailey, J.K. and Friars, G.W. 1994. Sources of variation in age at smolting in hatchery-reared Atlantic Salmon (*Salmo salar*). Canadian Journal of Animal Science 74: 223-227

⁰⁴Bailey, J.K., Oliver, G. and Friars, G.W. 1993. Inheritance of furunculosis resistance in Atlantic salmon. Proceedings of the 10th annual meeting of the Aquaculture Association of Canada 93: 90-92.

Bakke, T.A., Jansen, P.A. and Hansen, L.P. 1990. Differences in the host resistance of Atlantic salmon, *Salmo salar*, stocks to the monogenean *Gyrodactylus salaricus* Malmberg, 1957. Journal of Fish Biology 37:557-587.

Bakke, T. A., Soleng, A., Harris, P.D. and Hansen, L.P. 1996. Gyrodactylid infections of pure Atlantic salmon, *Salmo salar* and brown trout *Salmo trutta* and their hybrid progeny. Parasitology 119: 467-481.

Bardon, A., Vandeputte, M., Dupont-Nivet, M. Chavanne, H., Haffray, P., Vergnet, A., Chatain, B. 2009. What is the heritable component of spinal deformities in the European sea bass (*Dicentrarchus labrax*)? Aquaculture 294(3-4): 194-201.

⁰⁵Basiao, Z. U. and Doyle, R.W. 1999. Test of size-specific mass selection for Nile tilapia, *Oreochromis niloticus* L., cage farming in the Philippines. Aquaculture Research 30: 373-378.

⁰⁶Beacham, T.D. and Evelyn, T.P.T. 1992. Population and genetic variation in resistance of chinook salmon to vibriosis, furunculosis, and bacterial kidney disease. Journal of Aquatic Animal Health 4: 153-167.

Bell, A.E. Burris, M.J., 1973. Simultaneous selection for two correlated traits in *Tribulium*. Genetics Research, Cambridge 21:29-46.

⁰⁷Bongers, A.B.J.; Bovenhuis, H.; Vanstokkom, A.C.; Wiegertjes, G.F.; Zandiehoulabi, B.; Komen, J., and Richter, C.J. J. 1997. Distribution of genetic variance in gynogenetic or androgenetic families. Aquaculture 153: 225-238.

⁰⁸Brzeski, V.J. and Doyle, R.W. 1995. A test of an on-farm selection procedure for Tilapia growth in Indonesia. Aquaculture 137: 219-230.

⁰⁹Burger, G. and Chevassus, B. 1987 Potential of selection for age at first sexual maturation in rainbow trout (*Salmo gairdneri* R.). Bulletin Français de la Pêche et de la Pisciculture. Paris 307:102-117.

¹⁰Busack, C.A. and Gall, G.A.E. 1983. An initial description of the quantitative genetics of growth and reproduction in the mosquitofish, *Gambusia affinis*. Aquaculture 32: 123-140.

¹¹Chiayvareesajja, J.; Roed, K.H.; Eknath, A.E.; Danting, J.C.; de Vera, M.P., and Bentsen, H.B. 1999. Genetic variation in lytic activities of blood serum from Nile Tilapia and genetic associations with survival and body weight. *Aquaculture* 175: 49-62.

¹²Choe, M.K. and Yamazaki, F. 1998. Estimation of heritabilities of growth traits, and phenotypic and genetic correlations in juvenile masu salmon *Oncorhynchus masou*. *Fisheries Science* 64: 903-908.

¹³Clayton, G.M. 1990. Ichthyophthiriusis in fish: genetic variation in resistance to infection. *Dissertation Abstracts International B: Science & Engineering* 51: 241pp.

Clayton, G.A., Morris, J.A. and A. Robertson. 1957(a). An experimental check on quantitative genetical theory. 1. Short-term responses to selection. *Journal of Genetics* 55: 131-151.

Clayton G.A., Morris, J.A. and A. Robertson 1957(b). An experimental check on quantitative genetical theory II. The long-term effects of selection. *Journal of Genetics* 55: 152-170.

Conover, D.O., S.B. Munch and S.A. Arnott. 2009. Reversal of evolutionary downsizing caused by selective harvest of large fish. *Proc. R. Soc.* 176:2015-2020.

¹⁴Crandell, P.A. and Gall, G.A.E. 1993a. The genetics of age and weight at sexual maturity based on individually tagged rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 117: 95-105.

¹⁵Crandell, P.A. and Gall, G.A.E. 1993b. The effect of sex on heritability estimates of body weight determined from data on individually tagged rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 113: 47-55.

¹⁶Crandell, P.A. and Gall, G.A.E. 1993c. The genetics of body weight and its effect on early maturity based on individually tagged rainbow trout (*Oncorhynchus mykiss*) *Aquaculture* 117: 77-93.

Crozier, W.W. 1993. Evidence of genetic interaction between escaped farm salmon and wild Atlantic salmon (*Salmo salar* L.) in a Northern Irish river. *Aquaculture* 113: 19-29.

¹⁷Del Vallee, G., Taniguchi, N., Tsujimura, A. 1996. Genetic differences in some haematological traits of communally reared clonal ayu, *Plecoglossus altivelis* Temnick & Schlegel, under stressed and non-stressed conditions. *Aquaculture Research* 27: 787-793.

¹⁸Dorson, M., Quillet, E., Hollebecq, M.G., Torhy, C. and Chevassus, B. 1995. Selection of rainbow trout resistant to viral haemorrhagic septicaemia virus and transmission of resistance by gynogenesis. *Veterinary Research* 26: 361-368.

¹⁹Dunham, R.A. and Smitherman, R.O. 1983. Response to selection and realized heritability for body weight in three strains of channel catfish, *Ictalurus punctatus*, grown in earthen ponds. *Aquaculture* 33: 89-96.

- ²⁰Eknath, A.E. Dey, M.M., Rye, M., Gjerde, B., Abella, T.A., Sevilla, R., Tayamen, M.M., Reyes, R.A. and Bentsen, H.B. 1998. Selective breeding of Nile tilapia for Asia. Proceedings 6th World Congress on Genetics Applied to Livestock Production Armidale, Australia. Vol. 27: 89-96
- ²¹Elvingson, P. and Johansson, K. 1993. Genetic and environmental components of variation in body traits of rainbow trout (*Oncorhynchus mykiss*) in relation to age. *Aquaculture* 118: 191-204.
- Enfield, F.D. 1977 Selection experiments in *Tribolium* designed to look at gene action issues. In: Proceedings of the International Conference on Quantitative Genetics. Pp 177-190. Ed. E. Pollak, O. Kempthorne and T.B. Bailey Jr. The Iowa State University Press/Ames, Iowa, U.S.A.
- ²²Fevolden, S.E., Refstie, T. and Gjerde, B. 1993. Genetic and phenotypic parameters for cortisol and glucose stress response in Atlantic salmon and rainbow trout. *Aquaculture* 118: 205-216.
- ²³Fevolden, S.E., Roed, K.H. and Gjerde, B. 1994. Genetic components of post-stress and lysozyme activity in Atlantic salmon; correlations to disease resistance. *Fish & Shellfish Immunology* 7: 507-519.
- ²⁴Fevolden, S.E., Roed, K.H., Fjalestad, K.T. and Stien, J. 1999. Poststress levels of lysozyme and cortisol in adult rainbow trout: heritabilities and genetic correlations. *Journal of Fish Biology* 54: 900-910.
- Fisher, R.A. 1918. The correlation between relatives on the supposition of Mendelian inheritance. *Royal Society Edinburgh Transactions* 52: 399-433
- ²⁵Fjalestad, K.T. Larsen, H.J.S. and Roed, K.H. 1996. Antibody response in Atlantic salmon (*Salmo salar*) against *Vibrio anguillarum* and *Vibrio salmonicida* O-antigens: Heritabilities, genetic correlations and correlations with survival. *Aquaculture* 145: 77-90.
- Friars, G.W., Bohren, B.B. and McKean, H.E. 1962. Time trends in populations of chickens subjected to multiple objective selection. *Poultry Science* 41:1773-1784
- Friars, G.W., Chambers, J.R., Kennedy, A. and Smith, A.D. 1972. Selection for resistance to Marek's disease in conjunction with other economic traits in chickens. *Avian Diseases* 16: 2-10.
- Friars, G.W., Nayak, B.N. and Hurnik, J.F. 1973. Realized genetic parameters following selection in *Tribolium castaneum* in two environments. *Ann. Génét. Sélection Animal.* 5: 135-142.
- Friars, G.W., Bailey, J.K. and Saunders, R.L. 1979. Consideration of a method of analyzing diallel crosses of Atlantic salmon. *Canadian Journal of Genetics and Cytology* 21: 121-128.
- Friars, G.W., Bailey, J.K. and O'Flynn, F.M. 1990. An overview of selection practices in the Salmon Genetics Research Program. In: Canada-Norway finfish aquaculture workshop. Canadian Technical Report Fisheries and Aquatic Science 1761: 49-59.

²⁶Friars, G.W., Bailey, J.K. and O'Flynn, F.M. 1992. Merging conventional mating and selection practices with transgenic technology. Pp 39-42. In van der Meer Canadian Aquatic Biotechnology Network, National Research Council Institute for Marine Biosciences, Halifax N.S.

Friars, G.W., Bailey, J.K. and O'Flynn, F.M. 1993. Response to index selection for multiple traits in Atlantic salmon (*Salmo salar*). Aquaculture Association of Canada Bulletin 93-4: 41-43.

Friars, G.W., Bailey, J.K., O'Flynn, F.M., Carr, J.W., Anderson, J.M., McGreachy, S.A. and Bailey, T.J. 1994. A review of some genetic aspects of sea ranching and aquaculture in Atlantic salmon. Pp 56-63. In S. Calabi & A. Stout. A hard look at some tough issues. New England Atlantic salmon management conference. Salmon Association, Newport Maine.

Friars, G.W., Bailey, J.K. and O'Flynn, F.M. 1997. Realized genetic gains in economic traits of Atlantic salmon (*Salmo salar*). Aquaculture Association of Canada Special Publication 2: 69-73.

Friars, G.W., McMillan, I., Quinton, M.V., O'Flynn, F.M., McGeachy, S.A. and Benfey T.J. 2001. Family differences in relative growth of diploid and triploid Atlantic salmon (*Salmo salar* L.) Aquaculture 192: 23-29.

²⁷Fujio, Y., Nakajima, M., Nomura, Go. 1995. Selection response on thermal resistance of the guppy *Poecilia reticulata*. Fisheries Science 61: 731-734.

²⁸Gall, G.A.E. and Bakar, Y. 1999 Stocking density and tank size in the design of breed improvement programs for body size of tilapia. Aquaculture 173: 197-205.

Galton, F., 1897. The average contribution of each of several ancestors to the total heritage of the offspring. Proceeding of the Royal Society 61: 401-413.

Gardner, M.L. 1976. A review of factors which may influence the sea-age and maturation of Atlantic Salmon, *Salmo salar* L. Journal of Fish Biology 9: 289-327

Garenc, C. Silversides, F.G. and Guderly, H.. 1998. Boost swimming and its enzymatic correlates in the three spine stickle back. (*Gasterosteus aculeatus*): full-sib heritabilities. Canadian Journal of Zoology. 76: 680-688.

Gheyas, A.A., Woolliams, J.A., Taggart, J.B., Sattar, M.A., Das, T.K., McAndrew, B.J. Penman, D.J. 2009. Heritability estimation of silver carp (*Hypophthalmichthys molitrix*). Aquaculture 294(3-4):187-193.

Gjerde, B. 1988. Complete diallel crosses between six inbred groups of rainbow trout (*Salmo gairdneri*) Aquaculture 75: 71-87.

²⁹Gjedre, B. 1994. Response to individual selection for age at sexual maturity in Atlantic salmon, Aquaculture 38: 224-240.

Gjerde, B., Guanés, K. and Gjedrem, T. 1983. Effect of inbreeding on survival and growth in rainbow trout. *Aquaculture* 34: 327-332.

³⁰Gjedrem, T. 2004. Genetic improvement of cold-water fish species. *Aquaculture* 31(1): 25-33.

Gjedrem, T. Gjoen, H.M. and Gjerde, B. 1991. Genetic origin of Norwegian farmed salmon. *Aquaculture* 98: 41-50.

Gjedrem, T., Gjoen, H.M. and Gjerde, B. 1991a. Genetic origin of Norwegian farmed salmon. *Aquaculture* 98: 41-50.

³¹Gjedrem, T., Gjoen, H.M. and Salte, R. 1991b. Genetic variation in red cell membrane fragility in Atlantic salmon and rainbow trout. *Aquaculture* 98: 349-354.

³²Gjedrem, T., Salte, R., Gjoen, H. 1991c Genetic variation in susceptibility of Atlantic salmon to furunculosis. *Aquaculture* 97: 1-6.

Godo, O.R. 1995. Transplantation-tagging experiments in preliminary studies of migration of cod off Norway. *ICES Journal of Marine Science* 52: 955-962.

Guy, D.R., Bishop, S.C., Woolliams, J.A., Brotherstone, S. 2009. Genetic parameters for resistance to Infectious Pancreatic Necrosis in pedigreed Atlantic salmon (*Salmo salar*) post-smolts using a Reduced Animal Model. 2009. *Aquaculture* 290(3-4):229-235.

Hansen, M.H., Kenchington, E., Nielsen, E.E. 2001. Assigning individual fish to populations using microsatellite DNA markers. *Fish and Fisheries* 2: 93-112.

Hanke, A.R., Friars, G.W., Saunders, Q.L. and Terhune, J.M. 1989. Family x photo period interaction in juvenile Atlantic salmon, *Salmo salar*. *Genome* 32: 1105-1112.

³³Hard, J.J.; Connell, L.; Hershberger, W.K., and Harrell, L.W. 2000. Genetic variation in mortality of Chinook salmon during a bloom of the marine alga *Heterosigma akashiwo*. *Journal of Fish Biology* 56:1387-1397.

Hartman IV, J.L., Garvik, B. and L. Hartwell, 2001. Principles for the buffering of genetic variation. *Science* 291 (No.5506): 1001-1004.

Herbinger, C.M., and Friars, G.W. 1991. Correlation between condition factor and total lipid content in Atlantic salmon, *Salmo salar* L. parr. *Aquaculture and Fish Management* 22: 527-529.

Herbinger, C.M., Doyle, R.W., Pitman, E.R., Paquet, D., Mesa, K.A., Morris, D.B. Wright, J.M. and Cook, D. 1995. DNA fingerprint based analysis of paternal and maternal effects on offspring growth and survival in communally reared rainbow trout. *Aquaculture* 137: 245-256.

Hermida, M., Fernandez, C., Amaro, R., and Miguel, E.S. 2002. Heritability and “evolvability” of meristic characters in a natural population of *Gasterosteus aculeatus*. *Canadian Journal of Zoology* 80: 532-541.

³⁴Hershberger, W., Myers, J.M., Iwamoto, R.N., Macauley, N.C. and Saxton, A.M. 1990. Genetic changes in the growth of coho salmon (*Oncorhynchus kisutch*) in marine net-pens produced by ten years of selection. *Aquaculture* v85: 187-197.

³⁵Hetzel, D.J.S.; Crocos, P.J.; Davis, G.P.; Moore, S.S. and Preston, N.C. 2000. Response to selection and heritability for growth in the Kuruma prawn, *Penaeus japonicus*. *Aquaculture* 181: 215-224.

Hofer, A. 1998. Variance component estimation in animal breeding: a review. *Journal of Animal Breeding and Genetics* 115(4):247-265.

³⁶Horstgen-schwark, G. 1993. Selection experiments for improving “pan-size” body weight of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 112: 13-24.

³⁷Hulata, G. 1995. The history and current status of aquaculture genetics in Israel. *Israeli Journal of Aquaculture* 47: 142-154.

³⁸Huaag, C-M. and Liao, I-C. 1990. Response to mass selection for growth rate in *Oreochromis niloticus*. *Aquaculture* 85:199-205.

Hurley, S.M. and C.B. Schom, 1984. Genetic control of swimming stamina in Atlantic salmon (*Salmo salar*). *Canadian Journal of Genetics and Cytology*: 26: 57-61.

ICES Annual Science Conference. 2009. Berlin, Germany. Interactions between aquaculture and wild stocks: comparative experiences for Atlantic cod and Atlantic salmon. Website: <http://www.ices.dk/iceswork/asc/2009/themesessions.asp>. Theme Session Q.

³⁹Jarayabhand, P. and Thavornnyutikarn, M. 1995. Realized heritability estimation on growth rate of oyster, *Saccostrea cucullata* Born, 1778. *Aquaculture* 138: 111-118.

⁴⁰Jarimopas, P. 1986. Realized response of Thai red tilapia to weight-specific selection for growth. The First Asian Forum. Proceedings of the First Asian Fisheries Forum, Manila, Philippines, 26-31 May, pp 109-111.

⁴¹Jarimopas, P. 1990. Realized response to Thai red tilapia to 5 generations of size-specific selection for growth. The Second Asian Fisheries Forum. Proceedings of the Second Asian Fisheries Forum, Tokyo, Japan, 17-22 April 1989, pp 519-522.

Jonasson, J. 1996. Selection experiments on Atlantic salmon ranching 2. Variation among release sites and strains for return rate, body weight and ratio of grilse to total return. *Aquaculture* 144: 277-294.

- ⁴²Jonasson, J. 1997. Selection experiments in salmon ranching 1. Genetic and environmental sources of variation in survival and growth in fresh water. *Aquaculture* 157: 225-234.
- ⁴³Jonasson, J. and Gjedrem, T. 1993. Genetic and environmental sources of variation in survival and growth in freshwater. *Aquaculture* 109: 225-236.
- Jonasson, J. and Gjedrem, T. 1997. Genetic correlation for body weight of Atlantic salmon grilse between fish in sea ranching and land-based farming. *Aquaculture* 157: 205-214.
- Jonasson, J., B. Gjerde and T. Gjedrem. 1997. Genetic parameters for return rate and body weight of sea-ranched Atlantic salmon. *Aquaculture* 154: 219-231.
- Jui, P.Y. and Friars, G.W. 1974. Two-way selection on pupal weight with different degrees of inbreeding in *Tribolium castaneum*. *Canadian Journal of Genetics and Cytology* 16: 765-775.
- Jui, P.Y. and Friars, G.W. 1980. Performance of crosses of selected lines of *Tribolium castaneum*. *Canadian Journal of Genetics and Cytology* 22: 129-136
- Kanno, Y. and Kawamura, H. 1993a. Comparison of heritabilities estimated by two techniques of variance analyses for fry mortality rate of Masu salmon in Hokkaido. *Bulletin of the Japanese Society of Scientific Fisheries* 59: 1111-1116.
- ⁴⁴Kanno, Y. and Kawamura, H. 1993b. Estimations of heritability and inbreeding coefficient for early mortality rate in two strains of masu salmon in Hokkaido. *Bulletin of the Japanese Society of Scientific Fisheries* 59: 1117-1120.
- ⁴⁵Kamonrat, W. and Doyle, R.W. 1988. Genetic variation of scale circulus spacing (CIRC) in Tilapia hybrid (*Oreochromis mossambicus* x *O. urolepis hornorum*). The Second International Symposium on Tilapia in Aquaculture. Bangkok, Thailand, 16-20 March 1987, ICLARM Conference Proceedings. Manila, no. 15: 209-213.
- Khaw, H.L., Bovenhuis, H., Ponzoni, R.W., Rezk, M.A., Charo-Karisa, H., Komen, H. 2009. Genetic analysis of Nile tilapia (*Oreochromis niloticus*) selection line reared in two input environments. *Aquaculture* 294(1-2): 37-42.
- McKay, J. K.; Latta, R. G. 2002. Adaptive population divergence markers, QTL and traits. *Trends in Ecology and Evolution* 17: 285-291.
- ⁴⁶Knibb, W.; Gorshkova, G. and Gorshkov, S. 1997. Selection for growth in the gilthead seabream, *Sparus aurata* L. *Israeli Journal of Aquaculture* 49: 57-66.
- Law, R. and Grey, D.R. 1989. Evolution of yields from populations with age-specific cropping. *Evolutionary Ecology* 3: 343-359.
- ⁴⁷Lawrence, C.S. and Morrissy, N.M. 2000. Genetic improvement of marron *Cherax tenuimanus* Smith and yabbies *Cherax* spp. in Western Australia. *Aquaculture Research* 31: 69-82.

- Lerner, I.M. 1950. Population genetics and animal improvement. The Sydics of the Cambridge University Press, Bentley House, N.W.I., London. 342 pp.
- Lerner, I.M. 1958. The genetic basis of selection. John Wiley and Sons, Inc., New York. 298 pp.
- ⁴⁸McIntyre, J.D., Amend, D.F. 1978. Heritability of tolerance for infectious hematopoietic necrosis in sockeye salmon (*Oncorhynchus nerka*). Transactions of the American Fisheries Society 107: 305-308.
- McKay, L.R. 1982. A genetic analysis of growth in rainbow trout (*Salmo gairdneri*). Ph.D. thesis, University of Guelph, Guelph, Ontario Canada. xiii +200pp.
- ⁴⁹McKay, L.R., P.E. Ihssen and Friars, G.W. 1986. Genetic parameters of growth in rainbow trout, *Salmo gairdneri*, as a function of age and maturity. Aquaculture 58:241-254.
- McLeod, D.L. and Friars, G.W. 1986. A comparison of individual, family and index selection methods in *Tribolium castaneum* Canadian Journal of Genetics and Cytology 28: 301-305
- McMillan, I. and Robertson A.1974. The power of methods for the detection of major genes affecting quantitative characters. Heredity 32: 349-356.
- McMillan, I., Fairfull, R.W., Friars, G.W. and Quinton, M. 1995. The effect of simultaneous selection on the genetic correlation. Theoretical Applied Genetics 91:776-779
- ⁵⁰Meffe, GK, Weeks, S.C., Mulvey, M., Kandl, K.L. 1995. Genetic differences in thermal tolerance of eastern mosquitofish (*Gambusia holbrooki*; Poecillidae) from ambient and thermal ponds. Canadian Journal of Fisheries and Aquatic Sciences 52: 2704-2711.
- Merilä, J. and Sheldon, B. C. 2000. Lifetime reproductive success and heritability in nature. The American Naturalist No.3, Vol. 155: 301-310.
- ⁵¹Mustafa, A. and Mackinnon, B.M. 1999. Genetic variation in susceptibility of Atlantic salmon to the sea louse *Caligus elongatus* Nordmann, 1832. Canadian Journal of Zoology 77: 1332-1335.
- ⁵²Muruta, O., Harada, T., Miyashita, S., Izumi, K-I., Maeda, S., Kato, K., and Hidemi, K.1996. Selective breeding for growth in red sea bream. Fisheries Science 62: 845-849.
- ⁵³Myers, J.N., Hershberger, W.K., Saxton, A.M., Iwamoto, R.N. 2001. Estimates of genetic and phenotypic parameters for length and weight of marine net-pen reared coho salmon (*Oncorhynchus kisutch* Walbaum). Aquaculture Research 32: 277-285.
- Navarro, A., Zamorano, M.J., Hildebrandt, S., Gines, R., Aguilera, C., Afonso, J.M. 2009. Estimates of heritabilities and genetic correlations for growth and carcass traits in gilthead seabream (*Sparus auratus* L.) under industrial conditions. Aquaculture 289(3-4): 225-230.

- ⁵⁴Nilsson, J. 1990. Heritability estimates of growth-related traits in Arctic charr (*Salvelinus alpinus*). *Aquaculture* 84: 211-217.
- ⁵⁵Nilsson, J. 1992. Genetic variation in resistance of Arctic char to fungal infection. *Journal of Aquatic Animal Health*. 4: 126-128.
- ⁵⁶O'Flynn, F.M., Friars, G.W. and Bailey, J.K. 1992. Development of a selection index to improve market value of cultured Atlantic salmon (*Salmo salar*). *Genome* 35: 304-310.
- Peichei, C.L., Nereng, K.S., Ohgi, K.A., Cole, B.L., Colosimo, P.F., Buerkle, C.A., Schluter, D., Kingsley, D.M. 2001. The genetic architecture of divergence between threespine stickleback species. *Nature* 414: 901-905.
- ⁵⁷Perez, J.E. and Alfonsi, C. 1999. Selection and realized heritability for growth in the scallop, *Euvola ziczac* (L.). *Aquaculture Research* 30: 211-214.
- ⁵⁸Perez-Rostro, C.I.; Ramirez, J.L. and Ibarra, A.M. 1999. Maternal and cage effects on genetic parameter estimation for Pacific white shrimp *Penaeus vannamei* Boone. *Aquaculture Research* 30: 681-693.
- ⁵⁹Pottinger, T.G. and Carrick, T.R. 1999. Modification of the plasma cortisol response to stress in rainbow trout by selective breeding. *General and Comparative Endocrinology* 16: 122-132.
- Ricker, W.E. 1981. Changes in the average size and age of Pacific salmon. *Canadian Journal of Fisheries and Aquatic Science* 38: 1636–1656.
- Righton, D., Metcalfe, J., Connolly, P. 2001. Different behaviour of North and Irish Sea cod. *Nature* 411: 156.
- Ritland, K., 1996. A marker-based method for inferences about quantitative inheritance in natural populations. *Evolution* 50: 1062-1073.
- Ritter, J.A. Farmer, G.J., Misra, R.K., Goff, T.R., Bailey, J.K. and Baum, E.T. 1986. Parental influences and smolt size and sex ratio effects on sea age at first maturity of Atlantic salmon (*Salmo salar*) p.30-38. In D.J. Meerburg [ed]. *Salmonid age at maturity*. Canadian Special Publication Fisheries and Aquatic Science 1989.
- Robertson, A. and Lerner, I.M. 1949. The heritability of all-or-non traits: viability in poultry. *Genetics* 34: 395-411.
- ⁶⁰Roed, K.H., Brun, E., Larsen, H.J., Refstie, T. 1990. The genetic influence on serum haemolytic activity in rainbow trout. *Aquaculture* 85: 109-117.
- Roff, D.A. 1992. *The evolution of life histories - theory and analysis*. Chapman & Hall, London. 535pp.

Rose, G.A. 1993. Cod spawning on a migration highway in the Northwest Atlantic. *Nature* 366: 458-461.

⁶¹Rye, M. and Lillevik, K.M. 1990. Survival in the early fresh-water period in Atlantic salmon (*Salmo salar*) and rainbow trout (*Salmo gairdneri*): Heritabilities for survival and genetic correlation between survival and growth. *Aquaculture* 85: 328-329.

Saddler, S.E., Friars, G.W. and P.E. Ihssen. 1986. The influence of temperature and genotype on growth rate of hatchery reared salmonids. *Canadian Journal Animal Science* 66: 559-606

⁶²Sadler, S.E.; McLeod, D.; McKay, L.R., and Moccia, R.D. 1992. Selection for early spawning and repeatability of spawn date in rainbow trout. *Aquaculture* 100:103.

Sakomoto, T.; Danzman, R.G.; Okamoto, N.; Ferguson, M.M.; Ihssen, P.E. 1999. Linkage analysis of quantitative trait loci associated with spawning time in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 173: 33-43.

Sakomoto, T.; Danzman, R.G.; Garbi, K.; Howard, P.; Ozaki, A.; Khoo, S.K.; Worman, R.A.; Okamoto, N.; Ferguson, M.M.; Holm, I.; Guyomard, R.; Hoyheim, B. 2000. A microsatellite linkage map of rainbow trout (*Oncorhynchus mykiss*) characterized by large sex-specific differences in recombination rates. *Genetics* 155: 1331-1345.

Schaeffer, L.R. 2010. Personal communication.

Siegel, S. 1956. *Nonparametric statistics for the behavioural sciences*. McGraw-Hill, New York. 312 pp.

⁶³Silverstein, J.T., Hershberger, W.K. 1992. Precocious maturation in coho salmon (*Oncorhynchus kisutch*): Estimation of heritability. *Bulletin of the Aquaculture Association of Canada*. St. Andrews 92: 34-36.

⁶⁴Silverstein, J.T.; Hershberger, W.K. 1994. Genetic parameters of size pre- and post-smoltification in coho salmon (*Oncorhynchus kisutch*) *Aquaculture* 128: 67-77.

Smith, P.J. 1994. Genetic diversity of marine fisheries resources: possible impacts of fishing. *FAO Fisheries Technical Paper* 344. Rome, FAO 53p.

Smitherman, R.O., Dunham, R.A. and Whitehead, P.K. 1996. Selection hybridization on genome manipulation in Siluroidei. In the biology and culture of catfish. M. Legendre and J.P. Proteau eds. *Aquatic Living Resources* 9: Hors série, 93-102.

⁶⁵Smoker, W.W., Gharrett, A.J., Stekoll, M.S., Joyce, J.E. 1994. Genetic analysis of size in an anadromous population of pink salmon. *Canadian Journal of Fisheries and Aquatic Sciences* 51 (Suppl. 1): 9-15.

⁶⁶Standal, M., Gjerde, B. 1987. Genetic variation in survival of Atlantic salmon during the sea-rearing period. *Aquaculture*. 66:197-207.

Stearns, S.C. 1992. *The evolution of life histories*. Oxford University Press, Walton Street, Oxford, England. 244pp.

⁶⁷Stromsheim, A., Eide, D.M., Hofgaard, P.O., Larsen, H.J.S., Refsiter, T. Roed, K.H. 1994. Genetic variation in the humoral immune response against *Vibrio salmonicida* and in antibody titre against *Vibrio anguillarum* and total IgM in Atlantic salmon (*Salmo salar*). *Veterinary Immunology and Immunopathology* 44: 85-95.

⁶⁸Su, G.S.; Liljedahl, L.E., and Gall, G.A.E. 1997. Genetic and environmental variation of female reproductive traits in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 154: 115-124.

⁶⁹Su, G.S.; Liljedahl, L.E., and Gall, G.A. E. 1999. Estimates of phenotypic and genetic parameters for within-season date and age at spawning of female rainbow trout. *Aquaculture* 171: 209-220.

⁷⁰Sylven, S. and Elvingson, P.1992. Comparison of rainbow trout (*Oncorhynchus mykiss*) strains for body weight, length and age at maturity in different Swedish production systems. *Aquaculture* 104: 37-50.

⁷¹Taniguchi, N., Yamasak, M. and Tsujimura, A. 1996. Genetic and environmental variances of body size and morphological traits in communally reared clonal lines from gynogenetic diploid ayu *Plecoglossus altive*. *Aquaculture* 140: 333-341.

⁷²Tauk, M.T., Vermuelen, K., Bovenhuos, H., Kohen, H. 2001. Heredity of stress-related cortisol response in common carp (*Cyprinus carpio* L.) *Aquaculture* 199: 283-294.

⁷³Tave, D. 1994. Response to selection and realized heritability for length in golden shiner, *Notemigonus crysoleucas*. *Journal of Applied Aquaculture*. 4: 55-64.

Taylor, R.S., Kube, P.D., Muller, W.J., Elliott, N.G. 2009. Genetic variation of gross gill pathology and survival of Atlantic salmon (*Salmo salar* L.) during natural amoebic gill disease challenge. *Aquaculture* 294(3-4): 172-179.

⁷⁴Teichert-Coddington, D.R. Smitherman, R.O. 1988. Lack of response by *Tilapia nilotica* to mass selection for rapid early growth. *Transactions of the American Fisheries Society* 117: 297-300.

Tsujimura, A. and N. Taniguchi. 1996, Reduction of phenotypic variation in reproductive traits of clonal strains of Ayu *Plecoglossus altivalis*. *Nippon Suisan Gakkaishi* 62: 393-398.

Vandeputte, M. Dupont-Nivet, M., Chatin, B., Chevassus, B. 2001. Setting up a strain testing design for the sea bass *Dicentrarchus labrax*: a simulation study. *Aquaculture* 202: 329-342.

⁷⁵Wangila, B.C.C. and Dick, T.A. 1996. Genetic effects and growth performance in pure and hybrid strains of rainbow trout, *Oncorhynchus mykiss* (Walbaum) (order: Salmoniformes, family: Salmonidae). *Aquaculture research* 27: 35-41.

Wearden, 1979. Evolution of a philosophy of science for quantitative genetics. In *Variance components and animal breeding*. Eds. D. Van Vleck & S.R. Searle. Proceedings of a conference in honour of C.R. Henderson, July 16-17, 1979. Cornell University, New York. pp 3-18.

⁷⁶Wiegertjes, G.F., Stet, R.J.M., van Muiswinkel, W.B. 1995. Divergent selection for antibody production to produce standard carp (*Cyprinus carpio* L.) lines for the study of disease resistance in fish. *Aquaculture* 137: 257-262.

⁷⁷Wild, V., Simianer, H., Gjoen, H-M., Gjerde, B. 1994. Genetic parameters and genotype x environment interaction for early sexual maturity in Atlantic salmon (*Salmo salar*). *Aquaculture* 128: 51-65.

⁷⁸Winkelman, A.M. and Peterson, R.G. 1994a. Heritabilities, dominance variation, common environmental effects and genotype by environment interactions for weight and length in chinook salmon. *Aquaculture* 125: 17-30.

⁷⁹Winkelman, A.M. and Peterson, R.G. 1994b. Genetic parameters (heritabilities, dominance ratios and genetic correlations) for body weight and length of chinook salmon after 9 and 22 months of saltwater rearing. *Aquaculture* 125: 31-36.

⁸⁰Withler, R.E. Clarke, W.C., Riddell, B.E. Kreiberg, H. 1987. Genetic variation in freshwater survival and growth of Chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* 64: 85-96.

Wright, S. 1921. Systems of mating. *Genetics*, 6: 111-178.

Yamada, Y. 1962. Genotype by environment interaction and genetic correlation of the same trait under different environments. *Japanese Journal of Genetics*. 36: 498-509.

Yule, G.V. 1907. On the theory of inheritance of quantitative compound characters on the basis of Mendel's laws – A preliminary note. Report of the Third International Congress of Genetics. The Royal Horticultural Society, London. pp140-142.

⁸¹Zyczynski, A., Olech, W. Dobosz, K. Goryczki, K. and Kohlmann, K. 1995. An attempt to evaluate the genetic parameters of a rainbow trout (*Oncorhynchus mykiss*) population. *Fortschritte des Fischereiwissenschaft* 12: 77-83.

Correct citation for this publication:

Friars, G.W. and P. J. Smith. 2010. Heritability, correlation and selection response estimates of some traits in fish populations. Atlantic Salmon Federation Technical Report March 2010.

<http://asf.ca/docs/uploads/friars-smith.pdf> and

<http://www.huntsmanmarine.ca/html/publications.html>